THE JOURNAL OF CHEMICAL THERMODYNAMICS

Volume 20, 1988

EDITORS

M. L. McGlashan
Department of Chemistry
University College London
20 Gordon Street
London WC1H 0AJ
U.K.

P. A. G. O'Hare
Argonne National Laboratory
Chemical Technology Division
Argonne
Illinois 60439
U.S.A.

Assistant Editor: **M. B. Ewing** (University College London, U.K.)

ADVISORY BOARD

Chairman: G. M. Schneider (F.R.G.)

J. C. G. Calado (Portugal)
H. Chihara (Japan)
E. H. P. Cordfunke (The Netherlands)
J. Fuger (Commission of the European Communities)
M. Gaune-Escard (France)
C. Jolicoeur (Canada
K. N. Marsh (U.S.A.)
J. B. Ott (U.S.A.)
G. Pilcher (U.K.)
G. Somsen (The Netherlands)
F. L. Swinton (U.K.)
W. Wagner (F.R.G.)
R. D. Weir (Canada)
R. H. Wood (U.S.A)
W. A. Wóycicki (Poland)

ACADEMIC PRESS
(Harcourt Brace Jovanovich, Publishers)
London San Diego New York
Boston Sydney Tokyo Toronto

Copyright © 1988 Academic Press Limited

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 20

Number 1, January 1988

RÜBESAMEN, J.; SCHNEIDER, G. M. Excess volumes of liquid mixtures at high pressures. VI. Results for $\{(1-x)C_6H_6+xc\text{-}C_6H_{12}\}$ and $[(1-x)\{(C_2H_5)_3N$ or $(C_2H_5)_2O\}+xCH_3CCl_3]$ up to 30 and 150 MPa respectively. Pressure dependence of excess molar Gibbs energies 1

CONWAY, B. E.; AYRANCI, E. Structural effects in the partial molar volumes and isentropic compressibilities of organic bases and their conjugate ions . 9

ŁUSZCZYK, M. Volumetric behaviour of (3-methoxypropionitrile + water) in the vicinity of the upper critical solution temperature 29

LETCHER, T. M.; BAXTER, R. C. Excess enthalpies and excess volumes of (a bicyclic compound + 1-hexene or 1-hexyne or 1-heptene or 1-heptyne) at 298.15 K 39

EWING, M. B.; JOHNSON, K. A.; MCGLASHAN, M. L. The (liquid + liquid) critical state of (cyclohexane + methanol). IV. $(T, x)_p$ coexistence curve and the slope of the critical line 49

ROY, R. N.; HUFFORD, K.; LORD, P. J.; MRAD, D. R.; ROY, L. N.; JOHNSON, D. A. The first acidity constant of carbon dioxide in solutions of ammonium chloride from e.m.f. measurements at 278.15 to 318.15 K 63

RIBEIRO DA SILVA, M. A. V.; FERRÃO, M. L. C. C. H. Standard enthalpies of formation of tris(β-diketonate) iron(III) complexes: the mean (Fe–O) bond-dissociation enthalpies 79

NAGATA, I.; TAMURA, K. Excess molar enthalpies of (propan-1-ol or propan-2-ol + acetonitrile), (propan-1-ol or propan-2-ol + chlorobenzene), and (propan-1-ol or propan-2-ol + acetonitrile + chlorobenzene) at 298.15 K . 87

RIBEIRO DA SILVA, M. A. V.; PILCHER, G.; IRVING, R. J. Standard enthalpies of formation of the bis(pentane-2,4-dionato) complexes of the Group II metals: Be, Mg, Ca, Sr, Ba, Zn, Cd, and Hg. The mean (metal–oxygen) bond-dissociation enthalpies 95

KONINGS, R. J. M.; CORDFUNKE, E. H. P. The vapour pressures of hydroxides. I. The alkali hydroxides KOH and CsOH 103

FLANIGAN, D. A.; YESAVAGE, V. F.; CERISE, K. S.; NIESEN, V. G.; KIDNAY, A. J. Enthalpy of quinoline between 291 and 655 K at pressures to 10342 kPa . 109

LÜ HUI; FU HAIZHANG; JIA YONZHEN. Determination of activity coefficients at infinite dilution for each of 14 organic solutes in each of 3 organic solvents at 333.15 K by gas chromatography 119

ADVICE TO AUTHORS 125

Number 2, February 1988

BERNARD, C.; CHATILLON, C.; AIT-HOU, A.; HILLEL, R.; MONTEIL, Y.; BOUIX, J. Thermodynamics of (gallium + chlorine)(g). I. Vapour-pressure measurements and thermodynamic stability of GaCl(g), $GaCl_2$(g), $GaCl_3$(g), Ga_2Cl_2(g), Ga_2Cl_4(g), and Ga_2Cl_6(g) 129

KOWALSKA, M.; SKUDLARSKI, K.; BOTOR, J. The thermodynamic activity of SiO_2 in $\{x_1Cu_2O + x_2Na_2O + (1-x_1-x_2)SiO_2\}$(l) at 1373 K. 143

LETCHER, T. M.; BAXTER, R. C. Excess enthalpies and excess volumes of (a bicyclic compound + cyclohexene or 1,3-cyclohexadiene or 1,4-cyclohexadiene) at 298.15 K 149

WHITE, D. E.; WOOD, R. H.; BIGGERSTAFF, D. R. Heat capacities of 0.0150 mol·kg^{-1} NaCl(aq) from 604 to 718 K at 32 MPa 159

FLANIGAN, D. A.; YESAVAGE, V. F. Enthalpy of (*m*-cresol + 1,2,3,4-tetrahydronaphthalene) between 291 and 655 K at pressures to 10342 kPa . . 169

MORIYOSHI, T.; AONO, T. Compressions of C_6 hydrocarbons from 298.15 to 313.15 K at pressures to 145 MPa 185

IÑARREA, J.; VALERO, J.; PÉREZ, P.; GRACIA, M.; GUTIÉRREZ LOSA, C. H_m^E and V_m^E of some (butanone or dipropylether + an alkanol) mixtures . . . 193

TAKAHASHI, Y.; NAKAMURA, J.-I.; SMITH, J. F. Laser-flash calorimetry. V. Effect of oxygen contamination on the heat capacity of vanadium from 80 to 850 K. 201

MESSERLY, J. F.; TODD, S. S.; FINKE, H. L.; GOOD, W. D.; GAMMON, B. E. Condensed-phase heat-capacity studies and derived thermodynamic properties for six cyclic nitrogen compounds 209

CORDRAY, D. R.; IZATT, R. M.; CHRISTENSEN, J. J.† The excess enthalpies of (carbon dioxide + cyclohexane) at 390.15, 413.15, 438.15, 498.15, and 508.15 K from 7.50 to 14.39 MPa 225

CORDFUNKE, E. H. P.; OUWELTJES, W. Standard enthalpies of formation of uranium compounds. XIV. BaU_2O_7 and $Ba_2U_2O_7$ 235

RUIZ, B.; ROYO, F. M.; OTIN, S. Excess molar enthalpies of (an ether + a chlorinated compound) 239

EWING, M. B.; GOODWIN, A. R. H.; MCGLASHAN, M. L.; TRUSLER, J. P. M. Thermophysical properties of alkanes from speeds of sound determined using a spherical resonator. 2. *n*-Butane 243

Number 3, March 1988

FLANIGAN, D. A.; JOYCE, T. P.; YESAVAGE, V. F. Enthalpy of (quinoline + 1,2,3,4-tetrahydronaphthalene) between 291 and 655 K at pressures to 10342 kPa. 257

LUO, B.; BENSON, G. C.; LU, B. C.-Y. Excess enthalpies for (diethyl ether + *n*-alkane) at 298.15 K 267

BRUNNER, E. Fluid mixtures at high pressures. VI. Phase separation and critical phenomena in 18 (*n*-alkane + ammonia) and 4 (*n*-alkane + methanol) mixtures 273

BONDARENKO, A. A.; KOROBOV, M. V.; MITKIN, V. N.; SIDOROV, L. N. Enthalpy of sublimation of platinum tetrafluoride. 299

GREENHILL, D. A.; HOURSTON, D. J.; STUBLEY, D. Excess molar enthalpies of some pairs of model monomers of polystyrene and of polyacrylic esters . 305

PANKAJAVALLI, R.; SREEDHARAN, O. M. Thermodynamic stabilities of Mo_4O_{11}(s) and MoO_3(l) from the oxide e.m.f. method 309

PRASAD, R.; IYER, V. S.; ZILEY SINGH; VENUGOPAL, V.; MOHAPATRA, S.; SOOD, D. D. Thermodynamic properties of $Cr_{1-x}Te$ for $x = 0.546$ and 0.526 . . 319

WORMALD, C. J.; EYEARS, J. M. Excess molar enthalpies and excess molar volumes of $\{xCO_2+(1-x)C_2H_6\}$ up to 308.4 K and 11.0 MPa. . . 323

SIPOWSKA, J. T.; WIECZOREK, S. A. Excess molar volumes of (2,3,3-trimethylbutan-2-ol + heptane) between 293.15 and 308.15 K 333

MCCONVILLE, G. T.; PAVESE, F. Physicochemical problems involved in measuring thermodynamic properties of normal and equilibrium deuterium at the triple point. 337

RIBEIRO DA SILVA, M. A. V.; FERRÃO, M. L. C. C. H. Standard enthalpies of formation of bis(trifluoroacetylacetonato)copper(II) and bis(hexafluoroacetylacetonato)copper(II): the mean (Cu–O) bond-dissociation enthalpies . 359

KODEJŠ, Z.; ŠPALKOVA, H.; SACCHETTO, G. A. Vapour pressures of very concentrated electrolyte solutions. IV. Measurements on $\{(1-x)(CH_3)_2SO+xKSCN\}$ by a dew-point apparatus 363

NAVARRO, R.; PUERTOLAS, J. A.; PALACIO, F.; GONZALEZ, D. Thermal properties of tetraethylammonium tetrachloroferrate(III) and tetrabromoferrate(III) . 373

Number 4, April 1988

GIERYCZ, P.; GREGOROWICZ, J.; MALANOWSKI, S. Vapour pressures and excess Gibbs energies of (butan-1-ol + *n*-octane or *n*-decane) at 373.15 and 383.15 K 385

IMAMURA, A.; MURATA, S.; SAKIYAMA, M. Combustion and sublimation calorimetric studies on acetylurea and trimethyl isocyanurate 389

BOMMEL, M. J. VAN; MILTENBURG, J. C. VAN; SCHUIJFF, A. Heat-capacity measurements and thermodynamic functions of 1,3,5-triazine and 1,3,5-trioxane 397

MAKHATADZE, G. I.; PRIVALOV, P. L. Partial specific heat capacity of benzene and of toluene in aqueous solution determined calorimetrically for a broad temperature range 405

CLARK, A. Q.; STEAD, K. (Vapour + liquid) phase equilibria of binary, ternary, and quaternary mixtures of CH_4, C_2H_6, C_3H_8, C_4H_{10}, and CO_2 . . 413

KABO, G. J.; YURSHA, I. A.; FRENKEL, M. L.; POLESHCHUK, P. A.; FEDOSEENKO, V. I.; LADUTKO, A. I. Thermodynamic properties of cyclohexanol and cyclohexanone 429

BRUNNER, E. Effect of impurities on the critical temperature T^c at the upper critical end point of (cyclohexane + methanol) 439

LÓPEZ, J. A.; PÉREZ, P.; GRACIA, M.; GUTIÉRREZ LOSA, C. G^E_m(298.15 K) of mixtures containing 1,2-dichloroethane 447

PRASAD, R.; SMRUTI MOHAPATRA; IYER, V. S.; ZILEY SINGH; VENUGOPAL, V.; SOOD, D. D. Thermodynamics of iron telluride ($Fe_{0.540}Te_{0.460}$) . . . 453

CZARNOTA, I. Heat capacity of 2-methylbutane at high pressures . . . 457

LI SHAOFENG; PILCHER, G. Enthalpy of formation of pyridine-*N*-oxide: the dissociation enthalpy of the (N–O) bond 463

NAGATA, I. Isothermal (vapour + liquid) equilibria of (methanol + toluene) and of (methanol + acetonitrile + toluene) 467

YAO, S.; KOZUKA, Z. The activity coefficients of oxygen in liquid (germanium + bismuth), (germanium + lead), and (germanium + indium) alloys . . 473

PIKKARAINEN, L. Excess enthalpies of (2,2,2-trifluoroethanol + 2-butanone or *N*-methylacetamide or *N*,*N*-dimethylacetamide) 481

MESSERLY, J. F.; FINKE, H. L.; GOOD, W. D.; GAMMON, B. E. Condensed-phase heat capacities and derived thermodynamic properties for 1,4-dimethylbenzene, 1,2-diphenylethane, and 2,3-dimethylnaphthalene 485

FAUX, P. W.; CHRISTENSEN, J. J.; IZATT, R. M.; PANDO, C.; RENUNCIO, J. A. R. The excess enthalpies of (carbon dioxide + *n*-hexane + toluene) at 413.15 K and 7.50 and 12.50 MPa 503

Number 5, May 1988

AMIRKHANOV, KH. I.; ABDULAGATOV, I. M.; ALIBEKOV, B. G.; STEPANOV, G. V.; BOUY, O. A. Equation of state and thermodynamic properties of propan-2-ol 513

BORSHCHEVSKII, A. YA.; BOLTALINA, O. V.; SOROKIN, I. D.; SIDOROV, L. N. Thermochemical quantities for gas-phase iron, uranium, and molybdenum fluorides, and their negative ions 523

MACIEL, M. R. W.; FRANCESCONI, A. Z. Excess Gibbs free energies of (*n*-hexane + propan-1-ol) at 338.15 and 348.15 K and of (*n*-hexane + propan-2-ol) at 323.15, 338.15, and 348.15 K 539

ŠVÁB, L.; PĚTROŠ, L.; HYNEK, V.; SVOBODA, V. A calorimeter for the determination of enthalpies of vaporization at high temperatures and pressures . 545

MORIYOSHI, T.; UOSAKI, Y.; MATSUURA, H.; NISHIMOTO, W. (Liquid + liquid) equilibria of (water + ethanol + *n*-hexane) from 0.1 to 200 MPa at 298.15 K 551

PANDO, C.; RENUNCIO, J. A. R.; FAUX, P. W.; CHRISTENSEN, J. J.; IZATT, R. M. The excess enthalpies of (carbon dioxide + *n*-hexane + toluene) at 358.15 K and 7.50 and 12.50 MPa 559

CORDFUNKE, E. H. P.; OUWELTJES, W.; PRINS, G. Standard enthalpies of formation of tellurium compounds. III. Cs_2TeO_3, $Cs_2Te_2O_5$, $Cs_2Te_4O_9$, and Cs_2TeO_4 569

COLOMINA, M.; JIMÉNEZ, P.; PÉREZ-OSSORIO, R.; ROUX, M. V.; TURRIÓN, C. Thermochemical properties of benzoic acid derivatives. XVI. Vapour pressures and enthalpies of sublimation and formation of the three tetramethylbenzoic acids and of pentamethylbenzoic acid. 575

LETCHER, T. M.; BAXTER, R. C.; BEAN, A.; SEWRY, J. D. Activity coefficients of the tetramethyl compounds of Group 14 elements in *n*-alkane solutions from g.l.c. measurements 581

MORIYOSHI, T.; UOSAKI, Y.; SAKAMOTO, T. (Liquid + liquid) equilibria of (water + ethanol + trichloromethane) from 0.1 to 200 MPa at 298.15 K . . 589

HOVEY, J. K.; HEPLER, L. G.; TREMAINE, P. R. Apparent molar heat capacities and apparent molar volumes of $Hg(ClO_4)_2$(aq) and $Pb(ClO_4)_2$(aq) at 298.15 K 595

DYMOND, J. H.; MALHOTRA, R.; ISDALE, J. D.; GLEN, N. F. (p, ρ, T) of *n*-heptane, toluene, and oct-1-ene in the range 298 to 373 K and 0.1 to 400 MPa and representation by the Tait equation 603

ELKABULE, A. S.; WHITMAN, D. L.; MILLER, R. C. Excess molar enthalpies of (toluene + *n*-heptane) and (toluene + methylcyclohexane) using a new variable-volume dilution calorimeter 615

KLEINRAHM, R.; DUSCHEK, W.; WAGNER, W.; JAESCHKE, M. Measurement and correlation of the (pressure, density, temperature) relation of methane in the temperature range from 273.15 K to 323.15 K at pressures up to 8 MPa . 621

TRTANJ, M. I.; MAKSIMOVIĆ, Z. B.; ČOMOR, J. J. Excess molar volumes and enthalpies of (tetranitromethane + propene oxide or hexane or trichloromethane or tetrachloromethane) 633

NOTE

SVOBODA, V.; VESELÝ, F.; HYNEK, V.; CHVÁTAL, Z. Molar heat capacities of 1,1,1-trichlorotrifluoroethane, 1,4-dibromo-2,3-dichlorohexafluorobutane, and 1,6-dibromo-2,3,5-trichlorononafluorohexane 639

Number 6, June 1988

MORRISON, J. A. Thermodynamics of solid methanes: retrospective and prospective. (The 35th Huffman Memorial Award Lecture of the Calorimetry Conference delivered on 28 July 1987 at its 42nd annual conference held at Boulder, Colorado, U.S.A.). 641

CORDRAY, D. R.; IZATT, R. M.; CHRISTENSEN, J. J.; OSCARSON, J. L. The excess enthalpies of (carbon dioxide + ethanol) at 308.15, 325.15, 373.15, 413.15, and 473.15 K from 5.00 to 14.91 MPa 655

POMIANEK, T. Influence of aluminium and copper, as well as of relative difference of ionic volumes on the activity of Pb in $\{(1-x_1-x_2)\text{Ni} + x_1\text{Pb} + x_2(\text{Al or Cu})\}$ for $x_1, x_2 \ll 1$ 665

ROGERS, D. W.; DEJROONGRUANG, K. Enthalpies of hydrogenation of the *n*-heptenes and the methylhexenes 675

O'HARE, P. A. G.; LEWIS, B. M.; PARKINSON, B. A. Standard molar enthalpy of formation by fluorine-combustion calorimetry of tungsten diselenide (WSe_2). Thermodynamics of the high-temperature vaporization of WSe_2. Revised value of the standard molar enthalpy of formation of molybdenite (MoS_2) 681

EASTEAL, A. J.; WOOLF, L. A. (p, V_m, T, x) measurements for $\{(1-x)H_2O + xCH_3CN\}$ in the range 278 to 323 K and 0.1 to 280 MPa. I. Experimental results, isothermal compressibilities, isobaric expansivities, and partial molar volumes 693

EASTEAL, A. J.; WOOLF, L. A. (p, V_m, T, x) measurements for $\{(1-x)H_2O + xCH_3CN\}$ in the range 278 to 323 K and 0.1 to 280 MPa. II. Thermodynamic excess properties 701

ALONSO, R.; GUERRERO, R.; CORRALES, J. A. Excess molar enthalpies of (methylcyclohexane + an alkanol) at 323.15 K. II. Results for 2-methylpropan-1-ol, butan-2-ol, and 2-methylpropan-2-ol 707

RONDININI, S.; CAVADORE, A.; LONGHI, P.; MUSSINI, T. Redetermination of the standard potential of the mercury(I) sulphate electrode in aqueous solutions at temperatures from 283 to 343 K 711

JACKOWSKI, A. W. Multiple phase equilibria in (*n*-hexane + 4-octyl-4′-cyanobiphenyl) 721

KURIYAMA, N.; SAKIYAMA, M. Standard enthalpy of formation of $[Ni(NH_3)_6]Cl_2$(cr) as determined by solution-reaction calorimetry . . 729

COONEY, A.; MORCOM, K. W. Thermodynamic behaviour of mixtures containing fluoroalcohols. I. (Water + 2,2,2-trifluoroethanol) 735

KIRKLIN, D. R.; DOMALSKI, E. S. Enthalpies of combustion of triphenylphosphine and triphenylphosphine oxide 743

BRYNDZIA, L. T.; KLEPPA, O. J. Standard molar enthalpies of formation of realgar (α-AsS) and orpiment (As_2S_3) by high-temperature direct-synthesis calorimetry. 755

NOTES

AMBROSE, D.; GHIASSEE, N. B. Vapour pressures, critical temperatures, and critical pressures of 2-chloro-1,1,2-trifluoroethyl difluoromethyl ether (enflurane) and of 1-chloro-2,2,2-trifluoroethyl difluoromethyl ether (isoflurane) 765

AMBROSE, D.; GHIASSEE, N. B.; TUCKERMAN, R. Vapour pressure and critical temperature and critical pressure of 4-methylpentan-2-one . . . 767

Number 7, July 1988

MALLIKA, C.; SREEDHARAN, O. M. Standard molar Gibbs energy of formation of $MoTe_2$ from e.m.f. measurements 769

KONINGS, R. J. M.; CORDFUNKE, E. H. P.; OUWELTJES, W. The standard enthalpies of formation of hydroxides. II. The alkali hydroxides CsOH and KOH 777

IYER, V. S.; VENUGOPAL, V.; SMRUTI MOHAPATRA; ZILEY SINGH; ROY, K. N.; PRASAD, R.; SOOD, D. D. Standard molar Gibbs free energy of formation of Na_2ZrO_3(s) 781

KSIĄŻCZAK, A.; PARCZEWSKA, B. Vapour pressures of binary three-phase (solid + liquid + vapour) mixtures. I. Melting temperatures of polymorphic phases 785

KSIĄŻCZAK, A. Vapour pressures of binary three-phase (solid + liquid + vapour) mixtures. II. Melting temperatures of 1-hexadecanol-containing solid phases 791

AL-BIZREH, N.; WORMALD, C. J.; YERLETT, T. K. Enthalpies of $\{x(CH_3)_2CO + (1-x)C_6H_{14}\}$ for $x = 0.25$, 0.50, and 0.75 at 510.2 K up to 9.11 MPa 797

KASPRZYCKA-GUTTMAN, T.; VERA, J. H. Excess enthalpies and excess volumes of (2,6-lutidine + an *n*-alkane) 805

TAKAGI, T.; TERANISHI, H. Ultrasonic speed in (benzonitrile + nitrobenzene) under high pressures 809

BURRIEL, R.; TO, M. (W.K.); ZAINEL, H.; WESTRUM, E. F., JR.; CORDFUNKE, E. H. P.; MUIS, R. P.; WIJBENGA, G. Thermodynamics of uranium intermetallic compounds. II. Heat capacity of UPd_3 from 8 to 850 K 815

LAM, A. C.; ARCHER, D. G. Enthalpy of dilution of dodecyl- and tetradecyltrimethylammonium bromides in aqueous sodium bromide at elevated temperatures 825

DIOT, M.; GERMAIN, P.; EL MAÏL, R. Thermal properties of $TlTeF_5$ from 5 to 300 K. Solid-to-solid transition 843

BROWN, R. J. C.; CALLANAN, J. E.; WEIR, R. D.; WESTRUM, E. F., JR. The thermodynamics of ammonium indates. I. The heat capacity of the ammonium pentachloroindate monohydrate salt $(NH_4)_2InCl_5 \cdot H_2O$ from 8.5 to 349 K 847

PIKKARAINEN, L. Excess enthalpies of (phenylmethanol or 2-phenylethanol + 2-butanone or *N*-methylacetamide or *N*,*N*-dimethylacetamide) . . . 855

SAITO, A.; TANAKA, R. Excess volumes and heat capacities of binary mixtures formed from cyclohexane, hexane, and heptane at 298.15 K . . . 859

CHRISTENSEN, J. J.; CORDRAY, D. R.; OSCARSON, J. L.; IZATT, R. M. The excess enthalpies of four (carbon dioxide + an alkanol) mixtures from 308.15 to 573.15 K at 7.50 to 12.50 MPa 867

CORDRAY, D. R.; CHRISTENSEN, J. J.; IZATT, R. M.; OSCARSON, J. L. The excess enthalpies of (carbon dioxide + toluene) at 390.15, 413.15, 470.15, and 508.15 K from 7.60 to 17.50 MPa 877

JOHNSON, K. A. An asymmetric potential to describe the excess enthalpies of binary liquid mixtures near their (liquid + liquid) critical states . . 889

Number 8, August 1988

PANDO, C.; RENUNCIO, J. A. R.; FAUX, P. W.; CHRISTENSEN, J. J.; IZATT, R. M. The excess enthalpies of (carbon dioxide + *n*-hexane + toluene) at 308.15 K and 7.50 and 12.50 MPa 897

MEN DIAN-YUAN; LU DIAN-ZHEN; LIANG CHUN-YU; YANG JIA-ZHEN. Thermodynamic properties of multicomponent electrolyte solutions in mixed solvents. 4. $\{HCl(m_A) + NaCl(m_B)\}$ in $\{(1-x)H_2O + xCH_2OHCHOHCH_3\}$ at 278.15 to 318.15 K 907

MORIYA, K.; MATSUO, T.; SUGA, H. Calorimetric and dielectric studies of phase transitions in ammonium nitrite. 913

CARSON, A. S.; JAMEA, E. H.; LAYE, P. G.; SPENCER, J. A. The enthalpies of formation of 1,1-diphenylstannacyclopentane and 1,1-diphenylstannacyclohexane and some associated tin–carbon bond-enthalpy terms 923

ALONSO, R.; GUERRERO, R.; CORRALES, J. A. Excess molar enthalpies of (methylcyclohexane + an alkanol) at 323.15 K. III. Results for pentan-2-ol, pentan-3-ol, 2-methylbutan-1-ol, 3-methylbutan-1-ol, and 2-methylbutan-2-ol . . 927

JIMÉNEZ, J.; VALERO, J.; GRACIA, M.; GUTIÉRREZ LOSA, C. H^E_m of (an *n*-alkane + a butanol isomer) 931

RANDZIO, S. L.; EATOUGH, D. J.; LEWIS, E. A.; HANSEN, L. D. An automated calorimeter for the measurement of isobaric expansivities and of isothermal compressibilities of liquids by scanning pressure from 0.1 to 400 MPa at temperatures between 303 and 503 K. 937

MAIER, V.; GATES, J. A.; INGLESE, A.; WOOD, R. H. Volumetric properties of aqueous NaCl solutions from 0.0025 to 5.0 mol kg^{-1}, 323 to 600 K, and 0.1 to 40 MPa 949

RIBEIRO DA SILVA, M. A. V.; RIBEIRO DA SILVA, M. D. M. C.; PILCHER, G. Enthalpies of combustion of 1-hydroxynaphthalene, 2-hydroxynaphthalene, and 1,2-, 1,3-, 1,4-, and 2,3-dihydroxynaphthalenes 969

LUO WANG; BENSON, G. C.; LU, B. C. Y. Excess enthalpies for (di-*n*-propyl ether + *n*-alkane) at 298.15 K 975

VESELÝ, F.; ŠVÁB, L.; PROVAZNÍK, R.; SVOBODA, V. Enthalpies of vaporization at high pressures for methanol, ethanol, propan-1-ol, propan-2-ol, hexane, and cyclohexane 981

SKUDLARSKI, K.; DUDEK, J.; KAPALA, J. Thermodynamics of $\{xCdCl_2 + (1-x)CdBr_2\}$(s) investigated by mass spectrometry 985

KONINGS, R. J. M.; CORDFUNKE, E. H. P.; OUWELTJES, W. The standard enthalpies of formation of hydroxides. I. The alkaline-earth hydroxides β-$Ba(OH)_2$ and $Sr(OH)_2$ 989

AIT-HOU, A.; HILLEL, R.; CHATILLON, C. Mass-spectrometric study of gaseous (gallium + chlorine); molar enthalpies of formation of Ga_2Cl_4 and Ga_2Cl_6. 993

Number 9, September 1988

EUBANK, P. T.; JOFFRION, L. L.; PATEL, M. R.; WAROWNY, W. Experimental densities and virial coefficients for steam from 348 to 498 K with correction for adsorption effects 1009

Alikhanian, A. S.; Marushkin, K. N.; Greenberg, J. H.; Lazarev, V. B.; Gorgoraki, V. I. (p, T, x) phase diagram and thermodynamics of $\{(1-x)BaO+xWO_3\}$. 1035

Holmes, H. F.; Mesmer, R. E. Isopiestic studies of aqueous solutions at elevated temperatures. IX. $\{(1-y)NaCl+yLiCl\}$ 1049

Hossenlopp, I. A.; Archer, D. G. Enthalpies of vaporization of piperidine and 1,2-dimethylbenzene; gas-phase isobaric heat capacities of piperidine. . 1061

Duschek, W.; Kleinrahm, R.; Wagner, W.; Jaeschke, M. Measurement and correlation of the (pressure, density, temperature) relation of nitrogen in the temperature range from 273.15 to 323.15 K at pressures up to 8 MPa . 1069

Ott, J. B.; Purdy, J. E.; Neely, B. J.; Harris, R. A. Excess enthalpies at 323.15, 348.15, and 373.15 K, and (liquid + liquid) equilibria for (cyclohexane + acetonitrile) at 0.4 and 15 MPa 1079

Sun, T. F.; Schouten, J. A.; Trappaniers, N. J.; Biswas, S. N. Measurements of the densities of liquid benzene, cyclohexane, methanol, and ethanol as functions of temperature at 0.1 MPa 1089

García, I.; Cobos, J. C.; Escarda, R.; Casanova, C. Excess enthalpies of (butyric anhydride + *n*-alkane) at 298.15 K 1097

Nagata, I.; Tamura, K. Excess molar enthalpies of (butan-1-ol or 2-methylpropan-1-ol + acetonitrile), (2-methylpropan-1-ol + benzene), and (butan-1-ol or 2-methylpropan-1-ol + acetonitrile + benzene) at 298.15 K. . . 1101

Bender, T. M.; Van Hook, W. A. Molar volumes of CH_3OH, CH_3OD, and CD_3OD, and apparent or excess molar volumes for (methanol-OH + water), (methanol-OD + deuterium oxide), and (methanol-OH + methanol-OD), at 288.15, 298.15, and 313.15 K 1109

Notes

Letcher, T. M.; Sewry, J.; Orchard, S. W. Vapour pressure and density of tricyclo[3.3.0.$0^{2,6}$]octane 1115

Jain, V. K.; Whittenburg, S. L. Hypersonic speed, dynamic isentropic compressibility, and refractive indices for pyrimidine 1117

Book Review 1119

Number 10, October 1988

Haarhaus, U.; Schneider, G. M. (Liquid + liquid) phase equilibria in (methanol + butane) and (methanol + pentane) at pressures from 0.1 to 140 MPa 1121

KALINOWSKA, B.; WÓYCICKI, W. Heat capacities and excess heat capacities of (an alkanol + an unsaturated hydrocarbon). II. (Propan-1-ol + cyclohexene). 1131

LÓPEZ DE LA FUENTE, F. L.; CHEDA, J. A. R.; FERNÁNDEZ-MARTÍN, F.; WESTRUM, E. F., JR. Thermodynamics of thallium alkanoates. V. Heat capacity and thermodynamic functions of thallium(I) *n*-decanoate from 6 to 480 K . 1137

ASANO, M.; KOU, T. Thermochemical properties of $SrBO_2(g)$ and $SrB_2O_4(s)$. 1149

SAI BABA, M.; VISWANATHAN, R.; BALASUBRAMANIAN, R.; DARWIN ALBERT RAJ, D.; SAHA, B.; MATHEWS, C. K. Vaporization thermodynamics of (iron + tellurium): a high-temperature mass-spectrometric study 1157

FRENCH, H. T.; KOSHLA, M.; MARSH, K. N. Dielectric constants and apparent dipole moments of (butan-1-ol or butan-2-ol + cyclohexane) at 298.15 and 318.15 K and of (2-methylpropan-2-ol + cyclohexane) at 299.15 and 318.15 K 1175

SHIOHAMA, Y.; OGAWA, H.; MURAKAMI, S.; FUJIHARA, I. Excess molar isobaric heat capacities and isentropic compressibilities of (*cis*- or *trans*-decalin + benzene or toluene or iso-octane or *n*-heptane) at 298.15 K . . . 1183

BUGAJEWSKI, Z.; BYLICKI, A. (Solid + liquid) phase equilibria for (*N*-methylpyrrolidin-2-one + *m*-cresol) 1191

WANG XIANG-YUN; JIN TIAN ZHU; GOUDIAKAS, J.; FUGER, J. Thermodynamics of lanthanide elements. IV. Molar enthalpies of formation of $Y^{3+}(aq)$, $YCl_3(cr)$, $YBr_3(cr)$, and $YI_3(cr)$ 1195

MURATA, S.; SAKIYAMA, M.; SEKI, S. Standard enthalpies of formation of beryllium acetylacetonate(cr) in its stable and metastable states, beryllium oxypropionate(cr), and beryllium oxynitrate(cr) 1203

NUÑEZ, L.; BARRAL, L.; PILCHER, G. Enthalpies of combustion of oxamic acid, oxamide, and dithiooxamide 1211

DYMOND, J. H.; MALHOTRA, R.; AWAN, M. A.; ISDALE, J. D.; GLEN, N. F. (p, ρ, T) of $\{0.5C_8H_{18} + 0.5(1\text{-}C_8H_{16})\}$, $\{0.5(CH_3)_2CHCH_2C(CH_3)_3 + 0.5(1\text{-}C_8H_{16})$ or $0.5C_8H_{18}\}$, and $\{0.333C_8H_{18} + 0.333(1\text{-}C_8H_{16}) + 0.333(CH_3)_2CHCH_2C(CH_3)_3\}$, in the range 298 to 373 K and 0.1 to 500 MPa 1217

CARSON, A. S.; JAMEA, E. H.; LAYE, P. G.; SPENCER, J. A. The enthalpies of formation of triphenyl vinyl germanium, triphenyl phenylethynyl germanium, bis(triphenyl germanium) oxide, and some associated Ge–C bond enthalpies 1223

NOTE

AMBROSE, D.; GHIASSEE, N. B. Vapour pressure and critical temperature and critical pressure of 2,2,4,4,6,8,8-heptamethylnonane 1231

Number 11, November 1988

STEELE, W. V.; ARCHER, D. G.; CHIRICO, R. D.; COLLIER, W. B.; HOSSENLOPP, I. A.; NGUYEN, A.; SMITH, N. K.; GAMMON, B. E. The thermodynamic properties of quinoline and isoquinoline. 1233

LETCHER, T. M.; TAYLOR, S.; BAXTER, R. Excess molar enthalpies of (a cycloalkane + an alk-1-yne) at 298.15 K 1265

TOPOR, L.; KLEPPA, O. J. Standard molar enthalpies of formation of IrMe (Me = Ti, Zr, or Hf) and of OsTi by high-temperature mixing calorimetry. 1271

VESELÝ, F.; ZÁBRANSKÝ, M.; BARCAL, P.; SVOBODA, V. Molar heat capacities of four liquid fluorinated and halogenated ethanes. 1283

O'HARE, P. A. G.; LEWIS, B. M.; NGUYEN, S. N. Thermochemistry of uranium compounds. XVII. Standard molar enthalpy of formation at 298.15 K of dehydrated schoepite $UO_3 \cdot 0.9H_2O$. Thermodynamics of (schoepite + dehydrated schoepite + water) 1287

FAUX, P. W.; CHRISTENSEN, J. J.; IZATT, R. M.; PANDO, C.; RENUNCIO, J. A. R. The excess enthalpies of (carbon dioxide + *n*-hexane + toluene) at 470.15 K and 7.50 and 12.50 MPa 1297

SHIOHAMA, Y.; OGAWA, H.; MURAKAMI, S.; FUJIHARA, I. Excess thermodynamic properties of (*cis*-decalin or *trans*-decalin + cyclohexane or methylcyclohexane or cyclooctane) at 298.15 K 1307

SARMIENTO, F.; LÓPEZ, M.; PAZ ANDRADE, M. I.; FERNÁNDEZ, J.; ORTEGA, J.; PEÑA, J. A. Excess molar enthalpies at 298.15 K of (an *n*-alkyl formate + an *n*-alkanol). III. $\{xHCO_2CH_3 + (1-x)C_nH_{2n+1}OH\}$, ($n$ = 3 to 10) . . 1315

ACOSTA, S. M.; WESTRUM, E. F., JR.; BROWN, R. J. C.; CALLANAN, J. E.; WEIR, R. D. The thermodynamics of ammonium indates. II. The molar heat capacity of the ammonium pentabromoindate monohydrate salt $(NH_4)_2InBr_5 \cdot H_2O$ from 7.8 to 348 K 1321

GOZZI, D.; CARNEVALE, G.; PETRUCCI, L.; PETRUCCI, M. Thermodynamic equilibrium at a graphite-to-zirconia electrolyte interface 1327

FUKAI, M.; MATSUO, T.; SUGA, H. A calorimetric study of the short hydrogen-bond system. Heat capacities of potassium hydrogen maleate and potassium hydrogen fumarate 1337

STOKES, R. H. Excess enthalpies of (butan-1-ol + methanol) at 278.15, 298.15, and 318.15 K 1349

BASTOS, M.; NILSSON, S.-O.; RIBEIRO DA SILVA, M. D. M. C.; RIBEIRO DA SILVA, M. A. V.; WADSÖ, I. Thermodynamic properties of glycerol. Enthalpies of combustion and vaporization and the heat capacity at 298.15 K. Enthalpies of solution in water at 288.15, 298.15, and 308.15 K 1353

Number 12, December 1988

GILL, S. J. Calorimetry of model compounds that mimic biological molecules. (The 36th Huffman Memorial Award Lecture of the Calorimetry Conference delivered on 18 August 1988 at its 43rd annual conference held at Bartlesville, OK, U.S.A.) 1361

LEBEDEV, B. V.; KULAGINA, T. G.; SMIRNOVA, N. N. Thermodynamics of 4-methylcyclohexene, glycollide, and 1,1,3,3,5,5-hexaethylcyclotrisiloxane from 13.4 to 400 K 1383

BRUNNER, E. Fluid mixtures at high pressures. VII. Phase separations and critical phenomena in 12 binary mixtures containing ammonia . . . 1397

ITSUKI, H.; YAMASHITA, K.; TERASAWA, S. Partial molar volumes at infinite dilution for 1-$C_{m-1}H_{2m-1}X$ as solute in the *n*-alkane C_nH_{2n+2} as solvent for X = Cl, Br, I, CN, and OH, and $n = 6$, 9, and 12 at 298.15 K . . . 1411

REY, F. J.; MARTIN-GIL, J. Excess molar enthalpies of (a ketone + di-*n*-butyl ether) at 298.15 K 1417

CARSON, A. S.; DYSON, J.; LAYE, P. G.; SPENCER, J. A. The enthalpy of formation of 1,1-diphenylgermanocyclopentane and an assessment of the strain in the ring system. 1423

SEMENIUK, B.; KASPRZYCKA-GUTTMAN, T. (Vapour + liquid) equilibria of (β- or γ-picoline + a C_6 to C_9 *n*-alkane) at 348.15 K 1427

ABELLO, L.; CHHOR, K.; POMMIER, C.; SOURISSEAU, C. Thermodynamic study on perovskite-type layer compounds: $[NH_3(CH_2)_nNH_3]MCl(cr)$ (M = Mn, Cd; $n = 3$, 5) at low temperatures 1433

DEFOORT, F.; CHATILLON, C.; BERNARD, C. Mass-spectrometric study of (indium + chlorine)(g). Enthalpies of formation of InCl(g), $In_2Cl_2(g)$, $In_2Cl_4(g)$, $InCl_3(g)$, and $In_2Cl_6(g)$ 1443

GARCÍA, I.; COBOS, J. C.; ESCARDA, R.; CASANOVA, C. Excess enthalpies of (heptanoic anhydride + *n*-alkane) at 298.15 K 1457

JUNQUERA, E.; TARDAJOS, G.; AICART, E. Speeds of sound and isentropic compressibilities of (cyclohexane + benzene) and (1-chlorobutane + *n*-hexane or *n*-heptane or *n*-octane or *n*-decane) at 298.15 K . . . 1461

COONEY, A.; MORCOM, K. W. Thermodynamic behaviour of mixtures containing fluoroalcohols. II. (2,2,2-Trifluoroethanol + benzene) 1469

INDEX TO VOLUME 20 1477